



Approximate fixed point theorems for Geraghty-contractions

S. A. M. Mohsenialhosseini ^{a,*} H. Mazaheri ^b

^a*Faculty of Mathematics, Vali-e-Asr University, Rafsanjan, Iran—Faculty of Mathematics, Yazd University, Yazd, Iran*

^b*Faculty of Mathematics, Yazd University, Yazd, Iran*

Received 22 April 2015; accepted 25 June 2016

Abstract

The purpose of this paper is to obtain necessary and sufficient conditions for existence approximate fixed point on Geraghty-contraction. In this paper, definitions of approximate -pair fixed point for two maps T_α, S_α , and their diameters are given in a metric space.

Key words: Approximate fixed point; Approximate-pair fixed point; Geraghty-contraction.

2010 AMS Mathematics Subject Classification : 54H25, 54E50, 54E99, 68Q55.

* Corresponding author's E-mail: mohsenhosseini@yazd.ac.ir; amah@vru.ac.ir

1 Introduction

In 1973, Geraghty [2] introduced the Geraghty-contraction and proved the fixed point property for it. In 2006, MăDălina Berinde [1] proved the approximate fixed point property for various types of well known generalized contractions on metric spaces.

In this paper, starting from the article of Zhang, Su, Cheng [3], we study Geraghty-contraction on partially ordered metric spaces, and we give some qualitative and quantitative results regarding approximate fixed points of such contraction mapping.

Throughout this article, we denote by Γ the functions $\beta : [0, \infty) \rightarrow [0, 1)$ satisfying the following condition:

$$\beta(t_n) \rightarrow 1 \Rightarrow t_n \rightarrow 0.$$

Definition 1.1 [2] *Let (X, d) be a metric space. A mapping $T : X \rightarrow X$ is said to be a Geraghty-contraction if there exists $\beta \in \Gamma$ such that for any $x, y \in X$,*

$$d(Tx, Ty) \leq \beta(d(x, y))d(x, y).$$

Theorem 1.1 [2] *Let (X, d) be a complete metric space, and let $T : X \rightarrow X$ be an operator. Suppose that there exists $\beta \in \Gamma$ such that for any $x, y \in X$,*

$$d(Tx, Ty) \leq \beta(d(x, y))d(x, y).$$

Then T has a unique fixed point.

In 2012, Caballero et al. considered another contraction condition also give a generalization of Theorem 1.2 by considering a non-self mapping, and they get the following theorems.

Definition 1.2 [4] *Let A, B be two nonempty subsets of a metric space (X, d) . A mapping $T : A \rightarrow B$ is said to be a Geraghty-contraction if there exists $\beta \in \Gamma$ such that for any $x, y \in A$,*

$$d(Tx, Ty) \leq \beta(d(x, y))d(x, y).$$

In [1], the author defined the approximate fixed point property for self mapping on metric spaces.

Definition 1.3 [1] Let (X, d) be a metric space, $\epsilon > 0$ and $T : X \rightarrow X$ be a map. Then $x_0 \in X$ is ϵ -fixed point for T if $d(Tx_0, x_0) < \epsilon$.

Definition 1.4 [1] In this paper we will denote the set of all ϵ - fixed points of T , for a given ϵ , by :

$$F_\epsilon(T) = \{x \in X \mid x \text{ is an } \epsilon - \text{fixed point of } T\}.$$

Definition 1.5 [1] Let (X, d) be a metric space and $T : X \rightarrow X$ be a map. Then T has the approximate fixed point property if

$$\forall \epsilon > 0, F_\epsilon(T) \neq \emptyset.$$

Definition 1.6 [5] Let $(X, \|\cdot\|)$ be a completely norm space and $T : X \rightarrow X$, and $T_\alpha : X \rightarrow X$ be a map as follow:

$$T_\alpha = \alpha I + (1 - \alpha)T, \quad 0 < \alpha < 1.$$

Then $x_0 \in X$ is ϵ -fixed point for T_α if $\|T_\alpha x_0 - x_0\| < \epsilon$.

Remark 1.1 [5] In this paper we will denote the set of all ϵ - fixed points of T_α , for a given ϵ , by :

$$F_\epsilon(T_\alpha) = \{x \in X \mid x \text{ is an } \epsilon - \text{fixed point of } T_\alpha\}.$$

2 ϵ - fixed point in Geraghty-contraction for T and T_α maps

In this section, we give some results on ϵ - fixed point in Geraghty-contraction and its diameter.

Theorem 2.1 Let (X, d) be a metric space and $T : X \rightarrow X$ be a map, $x_0 \in X$ and $\epsilon > 0$. If $d(T^n(x_0), T^{n+k}(x_0)) \rightarrow 0$ as $n \rightarrow \infty$ for some $k > 0$, then T^k has an ϵ - fixed point.

Proof: . Since $d(T^n(x_0), T^{n+k}(x_0)) \rightarrow 0$ as $n \rightarrow \infty$, $\epsilon > 0$

$$\exists n_0 > 0 \text{ s.t. } \forall n \geq n_0 \quad d(T^n(x_0), T^{n+k}(x_0)) < \epsilon.$$

Then

$$d(T^{n_0}(x_0), T^k(T^{n_0}(x_0))) < \epsilon,$$

therefore $T^{n_0}(x_0)$ is an ϵ - fixed point of T^k . ■

Theorem 2.2 Let (X, d) be a metric space and $T : X \rightarrow X$ a Geraghty-contraction map. Then:

$$\forall \epsilon > 0, F_\epsilon(T) \neq \emptyset.$$

Proof: Let $\epsilon > 0$, $x \in X$.

$$\begin{aligned} d(T^n(x), T^{n+1}(x)) &= d(T(T^{n-1}(x)), T(T^n(x))) \\ &\leq \beta(d(T^{n-1}(x), T^n(x)))d(T^{n-1}(x), T^n(x)) \\ &\leq \dots \\ &\leq (\beta(d(T^{n-1}(x), T^n(x))))^{n-1}d(T(x), T^2(x)) \\ &\leq (\beta(d(T^{n-1}(x), T^n(x))))^n d(x, Tx). \end{aligned}$$

But $\beta \in \Gamma$ Therefore

$$\lim_{n \rightarrow \infty} d(T^n(x), T^{n+1}(x)) = 0, \forall x \in X.$$

Now by Theorem 2.3 it follows that $F_\epsilon(T) \neq \emptyset, \forall \epsilon > 0$. ■

Theorem 2.3 Let (X, d) be a metric space and $T : X \rightarrow X$ a Geraghty-contraction map. If $F_\epsilon(T)$, the set of Approximate fixed point of T , is nonempty then the mapping

$$T_\alpha = \alpha I + (1 - \alpha)T, \quad 0 < \alpha < 1$$

satisfy in Geraghty-contraction and $F_\epsilon(T) = F_\epsilon(T_\alpha)$. Moreover $d(T_\alpha^n(x), T_\alpha^{n+k}(x)) \rightarrow 0$ as $n \rightarrow \infty$, for some $k > 0$, $\epsilon > 0$.

Proof: By the definition of $F_\epsilon(T)$, $F_\epsilon(T) = F_\epsilon(T_\alpha)$. Also, since T satisfy in Geraghty-contraction and I is identify function, it follows that T_α satisfy in Geraghty-contraction. Now, we prove $d(T_\alpha^n(x_0), T_\alpha^{n+k}(x_0)) \rightarrow 0$ as $n \rightarrow \infty$. Suppose $x \in X$ now, observe first that $d(T_\alpha x, T_\alpha^2 x) \leq \beta(d(x, T_\alpha x))d(x, T_\alpha x)$ and, by induction, that $d(T_\alpha^n x, T_\alpha^{n+1} x) \leq (\beta(d(x, T_\alpha x)))^n d(x, T_\alpha x)$. Thus, for any n and any $k > 0$, we have

$$\begin{aligned} d(T_\alpha^n(x), T_\alpha^{n+k}(x)) &\leq \sum_{i=n}^{n+k-1} d(T_\alpha^i(x), T_\alpha^{i+1}(x)) \\ &\leq ((\beta(d(x, T_\alpha x)))^n + \cdots + (\beta(d(x, T_\alpha x)))^{n+k-1})d(x, T_\alpha x) \\ &\leq \frac{(\beta(d(x, T_\alpha x)))^n}{1 - (\beta(d(x, T_\alpha x)))} d(x, T_\alpha x). \end{aligned}$$

But $\beta \in \Gamma$ Therefore $d(T_\alpha^n(x_0), T_\alpha^{n+k}(x_0)) \rightarrow 0$ as $n \rightarrow \infty$. ■

Corollary 2.1 Let (X, d) be a metric space and $T : X \rightarrow X$ a Geraghty-contraction map. If $F_\epsilon(T)$, the set of Approximate fixed point of T , is nonempty then the mapping

$$T_\alpha = \alpha I + (1 - \alpha)T, \quad 0 < \alpha < 1$$

satisfy in Geraghty-contraction and $F_\epsilon(T) = F_\epsilon(T_\alpha)$. Then:

$$\forall \epsilon > 0, F_\epsilon(T_\alpha) \neq \emptyset.$$

Proof: By Theorem 2.4 it follows that $F_\epsilon(T) \neq \emptyset, \forall \epsilon > 0$, Therefore

$$F_\epsilon(T_\alpha) \neq \emptyset, \forall \epsilon > 0.$$

■

Definition 2.1 Let $T : X \rightarrow X$, be a map and $\epsilon > 0$. We define diameter $F_\epsilon(T)$ by

$$\text{diam}(F_\epsilon(T)) = \sup\{d(x, y) : x, y \in F_\epsilon(T)\}.$$

Theorem 2.4 Let $T : X \rightarrow X$, and $\epsilon > 0$. If $T : X \rightarrow X$ a Geraghty-contraction map. Then

$$\text{diam}(F_\epsilon(T)) \leq \frac{2\epsilon}{1 - \beta(d(x, Tx))}, \quad \beta \in \Gamma.$$

Proof. If $x, y \in F_\epsilon(T)$, then

$$\begin{aligned} d(x, y) &\leq d(x, Tx) + d(Tx, Ty) + d(Ty, y) \\ &\leq \epsilon_1 + \beta(d(x, Tx))d(x, y) + \epsilon_2. \end{aligned}$$

put $\epsilon = \text{Max}\{\epsilon_1, \epsilon_2\}$, therefore $d(x, y) \leq \frac{2\epsilon}{1 - \beta(d(x, Tx))}$. Hence $\text{diam}(F_\epsilon(T)) \leq \frac{2\epsilon}{1 - \beta(d(x, Tx))}$. ■

Definition 2.2 Let $T : X \rightarrow X$ a map,

$$T_\alpha = \alpha I + (1 - \alpha)T, \quad 0 < \alpha < 1$$

a map and $\epsilon > 0$. We define diameter $F_\epsilon(T_\alpha)$ by

$$\text{diam}(F_\epsilon(T_\alpha)) = \sup\{d(x, y) : x, y \in F_\epsilon(T_\alpha)\}.$$

Theorem 2.5 Let $T : X \rightarrow X$, and $\epsilon > 0$. If $T : X \rightarrow X$ a Geraghty-contraction map and $T_\alpha : X \rightarrow X$ be a map as follow:

$$T_\alpha = \alpha I + (1 - \alpha)T, \quad 0 < \alpha < 1$$

Then

$$\text{diam}(F_\epsilon(T_\alpha)) \leq \frac{2\epsilon}{1 - \beta(d(x, T_\alpha x))}.$$

Proof. If $x, y \in F_\epsilon(T_\alpha)$, then

$$\begin{aligned} d(x, y) &\leq d(x, T_\alpha x) + d(T_\alpha x, T_\alpha y) + d(T_\alpha y, y) \\ &\leq \epsilon_1 + \beta(d(x, T_\alpha x))d(x, y) + \epsilon_2. \end{aligned}$$

put $\epsilon = \text{Max}\{\epsilon_1, \epsilon_2\}$, therefore $d(x, y) \leq \frac{2\epsilon}{1 - \beta(d(x, T_\alpha x))}$. Hence $\text{diam}(F_\epsilon(T_\alpha)) \leq \frac{2\epsilon}{1 - \beta(d(x, T_\alpha x))}$. ■

3 Approximate-pair fixed point and (T_α, S_α)

In this section we will consider the existence of approximate fixed points for two maps $T_\alpha : A \cup B \rightarrow A \cup B$, $S_\alpha : A \cup B \rightarrow A \cup B$, where

$$T_\alpha = \alpha I + (1 - \alpha)T, \quad S_\alpha = \alpha I + (1 - \alpha)S, \quad 0 < \alpha < 1,$$

and $T : A \cup B \rightarrow A \cup B$, $S : A \cup B \rightarrow A \cup B$.

In 2011, Mohsenalhosseini et al. considered the existence of approximate best proximity points for two maps $T : A \cup B \rightarrow A \cup B$, $S : A \cup B \rightarrow A \cup B$ and they get the following theorems.

Definition 3.1 [6] *Let A and B be nonempty subsets of a metric space (X, d) and $T : A \cup B \rightarrow A \cup B$, $S : A \cup B \rightarrow A \cup B$ be two maps such that $T(A) \subseteq B$, $S(B) \subseteq A$. A point (x, y) in $A \times B$ is said to be an approximate-pair fixed point for (T, S) in X , if there exists $\epsilon > 0$*

$$d(Tx, Sy) \leq d(A, B) + \epsilon.$$

We say that the pair (T, S) has the approximate-pair fixed property in X if

$$P_{(T,S)}^a(A, B) \neq \emptyset,$$

where

$$P_{(T,S)}^a(A, B) = \{(x, y) \in A \times B : d(Tx, Sy) \leq d(A, B) + \epsilon \text{ for some } \epsilon > 0\}.$$

Theorem 3.1 [6] *Let A and B be nonempty subsets of a metric space (X, d) and $T : A \cup B \rightarrow A \cup B$, $S : A \cup B \rightarrow A \cup B$ be two maps such that $T(A) \subseteq B$, $S(B) \subseteq A$. If, for every $(x, y) \in A \times B$,*

$$d(T^n(x), S^n(y)) \rightarrow d(A, B)$$

then (T, S) has the approximate-pair fixed property.

Definition 3.2 *Let A and B be nonempty subsets of a metric space (X, d) and $T_\alpha : A \cup B \rightarrow A \cup B$, $S_\alpha : A \cup B \rightarrow A \cup B$ be two Geraghty-contraction maps such that $T_\alpha(A) \subseteq B$, $S_\alpha(B) \subseteq A$. A point (x, y) in*

$A \times B$ is said to be an approximate-pair fixed point for (T_α, S_α) in X , if there exists $\epsilon > 0$

$$d(T_\alpha x, S_\alpha y) \leq d(A, B) + \epsilon.$$

We say that the pair (T_α, S_α) has the approximate-pair fixed property in X if

$$P_{(T_\alpha, S_\alpha)}^\epsilon(A, B) \neq \emptyset,$$

where

$$P_{(T_\alpha, S_\alpha)}^\epsilon(A, B) = \{(x, y) \in A \times B : d(T_\alpha x, S_\alpha y) \leq d(A, B) + \epsilon \text{ for some } \epsilon > 0\}.$$

Theorem 3.2 Let A and B be nonempty subsets of a metric space (X, d) and $T_\alpha : A \cup B \rightarrow A \cup B$, $S_\alpha : A \cup B \rightarrow A \cup B$ be two maps such that $T_\alpha(A) \subseteq B$, $S_\alpha(B) \subseteq A$. If, for every $(x, y) \in A \times B$,

$$d(T_\alpha^n(x), S_\alpha^n(y)) \rightarrow d(A, B)$$

then (T_α, S_α) has the approximate-pair fixed property .

Proof. For $\epsilon > 0$, Suppose $(x, y) \in A \times B$. Since

$$d(T_\alpha^n(x), S_\alpha^n(y)) \rightarrow d(A, B)$$

$$\exists n_0 > 0 \text{ s.t. } \forall n \geq n_0 : d(T_\alpha^n(x), S_\alpha^n(y)) < d(A, B) + \epsilon$$

Then $d(T_\alpha(T_\alpha^{n-1}(x)), S_\alpha(S_\alpha^{n-1}(y))) < d(A, B) + \epsilon$ for every $n \geq n_0$. Put $x_0 = T_\alpha^{n_0-1}(x)$ and $y_0 = S_\alpha^{n_0-1}(y)$. Hence $d(T_\alpha(x_0), S_\alpha(y_0)) \leq d(A, B) + \epsilon$ and $P_{(T_\alpha, S_\alpha)}^\epsilon(A, B) \neq \emptyset$. ■

Definition 3.3 Let $T_\alpha : A \cup B \rightarrow A \cup B$, $S_\alpha : A \cup B \rightarrow A \cup B$ be continuous maps such that $T_\alpha(A) \subseteq B$, $S_\alpha(B) \subseteq A$. We define diameter $P_{(T_\alpha, S_\alpha)}^\epsilon(A, B)$ by,

$$\text{diam}(P_{(T_\alpha, S_\alpha)}^\epsilon(A, B)) = \sup\{d(x, y) : d(T_\alpha x, S_\alpha y) \leq \epsilon + d(A, B) \text{ for some } \epsilon > 0\}.$$

Example 3.6. Suppose $A = \{(x, 0) : 0 \leq x \leq 1\}$, $B = \{(x, 1) : 0 \leq x \leq 1\}$, $T(x, 0) = T(x, 1) = (\frac{1}{2}, 1)$ and $S(x, 1) = S(x, 0) = (\frac{1}{2}, 0)$. Then $d(T(x, 0), S(y, 1)) = 1$ and $\text{diam}(P_{(T_\alpha, S_\alpha)}^\epsilon(A, B)) = \text{diam}(A \times B) = \sqrt{2}$.

Theorem 3.3 Let $T_\alpha : A \cup B \rightarrow A \cup B$, $S_\alpha : A \cup B \rightarrow A \cup B$ be continuous maps such that $T_\alpha(A) \subseteq B$, $S_\alpha(B) \subseteq A$. If, there exists a $k \in [0, 1]$,

$$d(x, T_\alpha x) + d(S_\alpha y, y) \leq kd(x, y).$$

Then

$$\text{diam}(P_{(T_\alpha, S_\alpha)}^\epsilon(A, B)) \leq \frac{\epsilon}{1-k} + \frac{d(A, B)}{1-k} \text{ for some } \epsilon > 0.$$

Proof. If $(x, y) \in P_{(T_\alpha, S_\alpha)}^\epsilon(A, B)$, then

$$\begin{aligned} d(x, y) &\leq d(x, T_\alpha x) + d(T_\alpha x, S_\alpha y) + d(S_\alpha y, y) \\ &\leq \epsilon + kd(x, y) + d(A, B). \end{aligned}$$

Therefore $d(x, y) \leq \frac{\epsilon}{1-k} + \frac{d(A, B)}{1-k}$. Then $\text{diam}(P_{(T_\alpha, S_\alpha)}^\epsilon(A, B)) \leq \frac{\epsilon}{1-k} + \frac{d(A, B)}{1-k}$. ■

Acknowledge

The authors are extremely grateful to the referees for their helpful suggestions for the improvement of the paper.

References

- [1] M. Berinde, "Approximate Fixed Point Theorems," *Mathematica*, vol. LI, no. 1, 2006.
- [2] M. Geraghty, "On contractive mappings," *Proc. Amer. Math. Soc.* 40 (1973) 604-608.
- [3] Jingling Zhang, Yongfu Su, Qingqing Cheng, "Best proximity point theorems for generalized contractions in partially ordered metric spaces," *Fixed Point Theory and Applications* 2013, 2013-200.

- [4] J. Caballero et al., A best proximity point theorem for Geraghty-contractions, *Fixed Point Theory and Applications* 2012, 2012-231.
- [5] S.A.M. Mohsenalhosseini, H. Mazaheri, Fixed Point For completely norm Space and map T_α , *Mathematica Moravica*. Vol. 16-2 (2012), 25-35.
- [6] S. A. M. Mohsenalhosseini, H. Mazaheri, M. A. Dehghan, Approximate Best Proximity Pairs in Metric Space, *Abstract and Applied Analysis*, Volume 2011, Article ID 596971, 9 pages.