



Mathematics Scientific Journal

Vol. 7, No. 2, (2012), 1-3



Some notes on the existence of an inequality in Banach algebra

M. Asadi ^{a,*}

^a*Department of Mathematics, Zanzan Branch, Islamic Azad University,
Zanzan, Iran.*

Received 12 March 2011; accepted 1 June 2011

Abstract

We shall prove an existence inequality for two maps on Banach algebra, with an example and in sequel we have some results on \mathbb{R} and \mathbb{R}^n spaces. This way can be applied for generalization of some subjects of mathematics in teaching which how we can extend a math problem to higher level.

Key words: Banach algebra; Norm inequality.

2010 AMS Mathematics Subject Classification : 32A65; 47A30.

* Corresponding Author. Tel: (+98-241-4221001-7); Fax: (+98-241-4220030);
Email address: masadi@azu.ac.ir (M. Asadi).

1 Introduction

We generalize a problem which was a simple existence inequality in [1, page 229] to Banach algebra ([2]) and in continue we obtain some results on \mathbb{R} and \mathbb{R}^n spaces.

2 Main results

Theorem 2.1 *Let E be a nonempty Banach algebra over \mathbb{C} , f and g two self-maps on E . If there exists $a \in E$ such that $\|a^2\| \geq 1$, then*

$$\|xy - f(x) - g(y)\| \geq \frac{1}{4},$$

for some $x, y \in E$.

Proof. Let $\|xy - f(x) - g(y)\| < \frac{1}{4}$ for all $x, y \in E$. So we have $\|f(0) + g(0)\| < \frac{1}{4}$, $\|f(a) + g(0)\| < \frac{1}{4}$, $\|f(0) + g(a)\| < \frac{1}{4}$. But

$$\begin{aligned} \|a^2 - f(a) - g(a)\| &\geq \|a^2\| - \|f(a) + g(a)\| \\ &> 1 - \|f(a) + g(0)\| - \|g(0) + f(0)\| \\ &\quad - \|f(0) + g(a)\| \\ &> \frac{1}{4}. \end{aligned}$$

□

Corollary 1 *If E be a nonempty unital Banach algebra over \mathbb{C} , f and g two self-maps on E , then*

$$\|xy - f(x) - g(y)\| \geq \frac{1}{4},$$

for some $x, y \in E$.

Proof. If $\mathbf{1}$ denotes the unit in the Banach algebra E , then $\|\mathbf{1}\| = 1$. □

Example 2.2 ([1]) Let f and g be real valued functions of a real variable. Therefore there exist $x, y \in [0, 1]$ such that $|xy - f(x) - g(y)| \geq \frac{1}{4}$.

In sequel, fixed $n \in \mathbb{N}$.

Corollary 2 If $f_1, \dots, f_n : \mathbb{R} \rightarrow \mathbb{R}$ be maps, then

$$\forall a_i \geq 0 \quad \exists x_i \in [0, a_i] \quad \left| x_1 \cdots x_n - \sum_{i=1}^n f_i(x_i) \right| \geq \frac{a_1 \cdots a_n}{2^n},$$

for all $i = 1, 2, \dots, n$.

Lemma 2.3 If $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$, $I = [0, a_1] \times \cdots \times [0, a_n]$, $J = [0, b_1] \times \cdots \times [0, b_n]$, $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$, then

$$\exists x \in I, y \in J \quad |x \cdot y - f(x) - g(y)| \geq \frac{a \cdot b}{4},$$

where $x \cdot y$ is inner product x and y .

Acknowledgments

The author is grateful to a referee for his/her careful reading and valuable comments and suggestions which led to the present form of the paper. The author would also like to thank *Professor S. Mansour Vaezpour* for his helpful advise to present this paper. This research has been supported by the Zanjan Branch, Islamic Azad University, Zanjan, Iran.

References

- [1] G. Klambauer, *Mathematical Analysis, Marcel Dekker, Inc., New York, 1975.*
- [2] R. G. Douglas, *Banach Algebra Techniques in Operator Theory*, Springer-Verlag, New York, 1998.