

# A Numerical Approach for Solving Forth Order Fuzzy Differential Equations Under Generalized Differentiability

E. Ahmady <sup>a,\*</sup>, N. Ahmady <sup>b</sup>

<sup>a</sup>*Department of Mathematics, Shahr-e-Qods Branch, Islamic Azad University, Tehran, Iran*

<sup>b</sup>*Department of Mathematics, Varamin-Pishva Branch, Islamic Azad University, Varamin, Iran*

Received 17 December 2015; accepted 27 March 2016

---

## Abstract

In this paper a numerical method for solving forth order fuzzy differential equations under generalized differentiability is proposed. This method is based on the interpolating a solution by piecewise polynomial of degree 8 in the range of solution . We investigate the existence and uniqueness of solutions. Finally a numerical example is presented to illustrate the accuracy of the new technique.

*Key words:* Fuzzy differential equations, Numerical Method, Generalized differentiability

*2010 AMS Mathematics Subject Classification :* 65L05, 34A07.

---

---

\* Corresponding author's E-mail: e.ahmadi@qodsiau.ac.ir(E. Ahmady)

## 1 Introduction

Fuzzy differential equations are a suitable tool to model problem in science and engineering. There are many idea to define a fuzzy derivative and in consequence, to study fuzzy differential equations. The first and most popular approach is using the Hukuhara differentiability for fuzzy valued function. Hukuhara differentiability has the drawback that the solution of fuzzy differential equations need to have increasing length of its support, so in order to overcome this weakness, Bede and Gal [12], introduced the strongly generalized differentiability of fuzzy valued function. This concept allows us to solve the above-mentioned shortcoming, also the strongly generalized derivative is defined for a larger class of fuzzy valued functions than the Hukuhara derivatives.

Higher-order fuzzy differential equations under generalized differentiability is presented by Khastan in [23]. Khastan proposed a analytic method to solve higher-order fuzzy differential equations based on the selection different type of derivatives, they obtained several solution to fuzzy initial value problem. In this paper a numerical method for forth order fuzzy differential equations is presented. The idea of this method is based on interpolating the solution by polynomial of degree 8 in the range of solution, the step size used is of length  $H = 4h$ . Also existence and uniqueness of the solutions are proved.

The paper is organized as follows: In section 2, some basic definitions are brought. A new method for solving forth order fuzzy differential equations, also the existence and uniqueness are introduced in section 3 and 4 . A numerical example is presented in section 5 and finally conclusion is drawn.

## 2 Notation and definitions

First notations which shall be used in this paper are introduced.

We denote by  $\mathbb{R}_{\mathcal{F}}$ , the set of fuzzy numbers, that is, normal, fuzzy convex, upper semi-continuous and compactly supported fuzzy sets which

are defined over the real line.

For  $0 < r \leq 1$ , set  $[u]^r = \left\{ t \in \mathbb{R} \mid u(t) \geq r \right\}$ , and  $[u]^0 = cl \left\{ t \in \mathbb{R} \mid u(t) > 0 \right\}$ .

We represent  $[u]^r = [u^-(r), u^+(r)]$ , so if  $u \in \mathbb{R}_{\mathcal{F}}$ , the  $r$ -level set  $[u]^r$  is a closed interval for all  $r \in [0, 1]$ . For arbitrary  $u, v \in \mathbb{R}_{\mathcal{F}}$  and  $k \in \mathbb{R}$ , the addition and scalar multiplication are defined by  $[u + v]^r = [u]^r + [v]^r$ ,  $[ku]^r = k[u]^r$  respectively.

A triangular fuzzy number is defined as a fuzzy set in  $\mathbb{R}_{\mathcal{F}}$ , that is specified by an ordered triple  $u = (a, b, c) \in \mathbb{R}^3$  with  $a \leq b \leq c$  such that  $u^-(r) = a + (b - a)r$  and  $u^+(r) = c - (c - b)r$  are the endpoints of  $r$ -level sets for all  $r \in [0, 1]$ .

**Definition 2.1** ([19]) *The Hausdorff distance between fuzzy numbers is given by  $D : \mathbb{R}_{\mathcal{F}} \times \mathbb{R}_{\mathcal{F}} \rightarrow \mathbb{R}^+ \cup \{0\}$ ,*

$$D(u, v) = \sup_{r \in [0, 1]} \max \left\{ |u^-(r) - v^-(r)|, |u^+(r) - v^+(r)| \right\}. \quad (2.1)$$

Consider  $u, v, w, z \in \mathbb{R}_{\mathcal{F}}$  and  $\lambda \in \mathbb{R}$ , then the following properties are well-known for metric  $D$ ,

1.  $D(u \oplus w, v \oplus w) = D(u, v)$ , for all  $u, v, w \in \mathbb{R}_{\mathcal{F}}$ ,
2.  $D(\lambda u, \lambda v) = |\lambda| D(u, v)$ , for all  $u, v \in \mathbb{R}_{\mathcal{F}}$ ,  $\lambda \in \mathbb{R}$
3.  $D(u \oplus v, w \oplus z) \leq D(u, w) + D(v, z)$ , for all  $u, v, w, z \in \mathbb{R}_{\mathcal{F}}$ ,
4.  $D(u \ominus v, w \ominus z) \leq D(u, w) + D(v, z)$ , as long as  $u \ominus v$  and  $w \ominus z$  exist, where  $u, v, w, z \in \mathbb{R}_{\mathcal{F}}$ .

where,  $\ominus$  is the Hukuhara difference ( $H$ -difference), it means that  $w \ominus v = u$  if and only if  $u \oplus v = w$ .

**Definition 2.2** ([12]) *Let  $u, v \in \mathbb{R}_{\mathcal{F}}$ . If there exists  $w \in \mathbb{R}_{\mathcal{F}}$  such that*

$$u \ominus_{gH} v = w \Leftrightarrow \begin{cases} (i) & u = v + w, \\ & \text{or} \\ (ii) & v = u + (-1)w, \end{cases}$$

Then  $w$  is called the generalized Hukuhara difference of  $u$  and  $v$ .

**Remark 2.1** Throughout the rest of this paper, we assume that  $u \ominus_{gH} v \in \mathbb{R}_{\mathcal{F}}$ .

Note that a function  $f : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}_{\mathcal{F}}$  is called fuzzy-valued function. The  $r$ -level representation of this function is given by  $f(t; r) = [f^-(t; r), f^+(t; r)]$ , for all  $t \in [a, b]$  and  $r \in [0, 1]$ .

**Definition 2.3** ([15]) A fuzzy valued function  $f : [a, b] \rightarrow \mathbb{R}_{\mathcal{F}}$  is said to be continuous at  $t_0 \in [a, b]$  if for each  $\epsilon > 0$  there is  $\delta > 0$  such that  $D(f(t), f(t_0)) < \epsilon$ , whenever  $t \in [a, b]$  and  $|t - t_0| < \delta$ . We say that  $f$  is fuzzy continuous on  $[a, b]$  if  $f$  is continuous at each  $t_0 \in [a, b]$ .

**Definition 2.4** ([15]) The generalized Hukuhara derivative of the fuzzy-valued function  $f : (a, b) \rightarrow \mathbb{R}_{\mathcal{F}}$  at  $t_0 \in (a, b)$  is defined as

$$f'_{gH}(t_0) = \lim_{h \rightarrow 0} \frac{f(t_0 + h) \ominus_{gH} f(t_0)}{h}. \quad (2.2)$$

If  $f'_{gH}(t_0) \in \mathbb{R}_{\mathcal{F}}$  satisfying (2.2) exists, we say that  $f$  is generalized Hukuhara differentiable ( $gH$ -differentiable for short) at  $t_0$ .

**Definition 2.5** ([15]) Let  $f : [a, b] \rightarrow \mathbb{R}_{\mathcal{F}}$  and  $t_0 \in (a, b)$ , with  $f^-(t; r)$  and  $f^+(t; r)$  both differentiable at  $t_0$  for all  $r \in [0, 1]$ . We say that

- $f$  is [(i) -  $gH$ ]-differentiable at  $t_0$  if

$$f'_{i.gH}(t_0; r) = [(f^-)'(t_0; r), (f^+)'(t_0; r)], \quad (2.3)$$

- $f$  is [(ii) -  $gH$ ]-differentiable at  $t_0$  if

$$f'_{ii.gH}(t_0; r) = [(f^+)'(t_0; r), (f^-)'(t_0; r)]. \quad (2.4)$$

**Definition 2.6** ([15]) We say that a point  $t_0 \in (a, b)$ , is a switching point for the differentiability of  $f$ , if in any neighborhood  $V$  of  $t_0$  there exist points  $t_1 < t_0 < t_2$  such that

**type(I)** at  $t_1$  (2.3) holds while (2.4) does not hold and at  $t_2$  (2.4) holds and (2.3) does not hold, or

**type(II)** at  $t_1$  (2.4) holds while (2.3) does not hold and at  $t_2$  (2.3) holds and (2.4) does not hold.

**Theorem 2.1** [9] Let  $T = [a, a+\beta] \subset \mathbb{R}$ , with  $\beta > 0$  and  $f \in \mathcal{C}_{gH}^n([a, b], \mathbb{R}_{\mathcal{F}})$ . For  $s \in T$

(i) If  $f^{(i)}$ ,  $i = 0, 1, \dots, n-1$  are  $[(i) - gH]$ -differentiable, provided that type of  $gH$ -differentiability has no change. Then

$$\begin{aligned} f(s) &= f(a) \oplus f'_{i.gH}(a) \odot (s-a) \oplus f''_{i.gH}(a) \odot \frac{(s-a)^2}{2!} \\ &\oplus \dots \oplus f^{(n-1)}_{i.gH}(a) \odot \frac{(s-a)^{n-1}}{(n-1)!} \oplus R_n(a, s), \end{aligned}$$

where

$$R_n(a, s) := \int_a^s \left( \int_a^{s_1} \dots \left( \int_a^{s_{n-1}} f^{(n)}_{i.gH}(s_n) ds_n \right) ds_{n-1} \dots \right) ds_1.$$

(ii) If  $f^{(i)}$ ,  $i = 0, 1, \dots, n-1$  is  $[(ii) - gH]$ -differentiable, provided that type of  $gH$ -differentiability has no change. Then

$$\begin{aligned} f(s) &= f(a) \ominus (-1)f'_{ii.gH}(a) \odot (s-a) \ominus (-1)f''_{ii.gH}(a) \\ &\odot \frac{(a-s)^2}{2!} \ominus (-1) \dots \ominus (-1)f^{(n-1)}_{ii.gH}(a) \\ &\odot \frac{(a-s)^{n-1}}{(n-1)!} \ominus (-1)R_n(a, s), \end{aligned}$$

where

$$R_n(a, s) := \int_a^s \left( \int_a^{s_1} \dots \left( \int_a^{s_{n-1}} f^{(n)}_{ii.gH}(s_n) ds_n \right) ds_{n-1} \dots \right) ds_1.$$

(iii) If  $f^{(i)}$  are  $[(i) - gH]$ -differentiable for  $i = 2k-1$ ,  $k \in \mathbb{N}$ , and  $f^{(i)}$  are  $[(ii) - gH]$ -differentiable for  $i = 2k$ ,  $k \in \mathbb{N} \cup \{0\}$ . Then

$$\begin{aligned}
f(s) &= f(a) \ominus (-1)f'_{ii.gH}(a) \odot (s-a) \oplus f''_{i.gH}(a) \\
&\odot \frac{(a-s)^2}{2!} \ominus (-1) \dots \ominus (-1)f^{(\frac{i-1}{2})}_{ii.gH}(a) \odot \frac{(a-s)^{\frac{i}{2}-1}}{(\frac{i}{2}-1)!} \\
&\oplus f^{(\frac{i}{2})}_{i.gH}(a) \odot \frac{(a-s)^{\frac{i}{2}}}{(\frac{i}{2})!} \ominus (-1) \dots \ominus (-1)R_n(a, s),
\end{aligned}$$

where

$$\begin{aligned}
R_n(a, s) &:= \\
&\int_a^s \left( \int_a^{s_1} \dots \left( \int_a^{s_{n-1}} f^{(n)}_{i.gH}(s_n) ds_n \right) ds_{n-1} \dots \right) ds_1.
\end{aligned}$$

(iv) Suppose that  $f \in \mathcal{C}_{gH}^n([a, b], \mathbb{R}_{\mathcal{F}})$ ,  $n \geq 3$ .

Furthermore let  $f$  in  $[a, \xi]$  is  $[(i) - gH]$ -differentiable and in  $[\xi, b]$  is  $[(ii) - gH]$ -differentiable, in fact  $\xi$  is switching point type I for first order derivative of  $f$  and  $t_0 \in [a, \xi]$  in a neighborhood of  $\xi$ . Moreover suppose that second order derivative of  $f$  in  $\zeta_1$  of  $[t_0, \xi]$  have switching point type II. Moreover type of differentiability for  $f^{(i)}$ ,  $i \leq n$  on  $[\xi, b]$  don't change. So

$$\begin{aligned}
f(s) &= f(t_0) \oplus f'_{i.gH}(t_0) \odot (\xi - t_0) \ominus f''_{ii.gH}(t_0) \\
&\odot (t_0 - \zeta_1) \odot (\xi - t_0) \oplus f''_{i.gH}(\zeta_1) \\
&\odot \left( \frac{(\xi - \zeta_1)^2}{2} - \frac{(t_0 - \zeta_1)^2}{2} \right) \ominus (-1)f'_{ii.gH}(\xi) \\
&\odot (s - \xi) \ominus (-1)f''_{ii.gH}(\xi) \odot \frac{(s - \xi)^2}{2!} \\
&\ominus (-1) \int_{t_0}^{\xi} \left( \int_{t_0}^{\zeta_1} \left( \int_{t_0}^{s_2} f'''_{ii.gH}(s_4) ds_4 \right) ds_2 \right) ds_1 \\
&\oplus \int_{t_0}^{\xi} \left( \int_{\zeta_1}^{s_1} \left( \int_{\zeta_1}^{s_3} f'''_{i.gH}(s_5) ds_5 \right) ds_3 \right) ds_1 \\
&\ominus (-1) \int_{\xi}^s \left( \int_{\xi}^{t_1} \left( \int_{t_0}^{t_2} f'''_{ii.gH}(t_3) dt_3 \right) dt_2 \right) dt_1.
\end{aligned}$$

### 3 Proposed Method

Consider the following fourth order fuzzy differential equation

$$\begin{cases} y^{(4)}(t) = f(t, y(t)), & t \in I = [0, T], \\ y(0) = y_0, y'(0) = y'_0, y''(0) = y''_0, y^{(3)}(0) = y_0^{(3)}, \end{cases} \quad (3.1)$$

where the derivative  $y^{(i)}$ ,  $i = 1, 2, 3, 4$ , is considered in the sense of gH-differentiability. The interval  $I$  may be  $[0, T]$  for some  $T > 0$  or  $I = [0, \infty)$ . We assume that  $f : I \times \mathbb{R}_{\mathcal{F}} \rightarrow \mathbb{R}_{\mathcal{F}}$  is sufficiently smooth function, and there exists  $k > 0$  such that

$$D(f(t, x), f(t, z)) \leq kD(x, z) \quad \forall t \in I, \quad x, z \in \mathbb{R}_{\mathcal{F}}. \quad (3.2)$$

Our construction of the fuzzy approximate solution  $s(t)$  is as follows: let  $y(t)$  be the fuzzy solution of (3.1), we divided the range of solution  $[0, T]$  into sub-intervals of equal length  $H = 4h = \frac{T}{n}$ , and let  $I_k = [kH, (k+1)H]$ , where  $k = 0, \dots, n-1$ . In this paper we approximate fuzzy solution of (3.1) by fuzzy piecewise polynomial of order 8. Piecewise approximation solution  $s(t)$  on  $I_k = [kH, (k+1)H]$ , is construct step by step as follows:

**Step One:** We define the first component of  $s(t)$  by  $s_0(t)$ , in two cases:

**Case(i):** Let us suppose that the unique solution of problem (3.1),  $y^{(i)}(t)$  are  $[(i) - gH]$ -differentiable, therefore  $s_0(t)$ , where in this cases is called  $s_{0,1}(t)$  for  $0 \leq t \leq H$  is as following

$$s_{0,1}(t) = \sum_{i=0}^4 y_{i.gH}^{(i)}(0) \odot \frac{t^i}{i!} \oplus \sum_{i=5}^8 \alpha_{i,0} \odot \frac{t^i}{i!}, \quad 0 \leq t \leq H, \quad (3.3)$$

**Case(ii):** Now, consider  $y^{(i)}(t)$  are  $[(ii) - gH]$ -differentiable, then  $s_0(t)$  that is called  $s_{0,2}(t)$  obtained for  $0 \leq t \leq H$  as follows:

$$s_{0,2}(t) = y(0) \ominus (-1) \sum_{i=1}^4 y_{ii.gH}^{(i)}(0) \odot \frac{t^i}{i!} \oplus \sum_{i=5}^8 \alpha_{i,0} \odot \frac{t^i}{i!}, \quad (3.4)$$

**Case(iii):** Now, consider  $y^{(i)}(t)$  are  $[(ii) - gH]$ -differentiable for  $i = 0, 2$ , and  $y^{(i)}(t)$  are  $[(i) - gH]$ -differentiable for  $i = 1, 3$ , then in this case  $s_0(t)$ , that is called  $s_{0,3}(t)$  is obtained for  $0 \leq t \leq H$  as follows:

$$\begin{aligned} s_{0,3}(t) = & y(0) \ominus (-1)y'_{ii.gH}(0) \odot t \oplus y''_{i.gH}(0) \odot \frac{t^2}{2} \\ & \ominus (-1)y^{(3)}_{ii.gH}(0) \odot \frac{t^3}{3!} \oplus y^{(4)}_{i.gH}(0) \odot \frac{t^4}{4!} \\ & \oplus \sum_{i=5}^8 \alpha_{i,0} \odot \frac{t^i}{i!}, \end{aligned} \quad (3.5)$$

In Eqs (3.3),(3.4) and (3.5), the coefficients  $\alpha_{i,0}$  for  $i = 5, 6, 7, 8$  as yet undetermined and to be obtained where  $s_0(t)$  satisfy the relations:

$$s_0^{(4)}(jh) = f(jh, s_0(jh)), \quad (3.6)$$

for  $j = 1, 2, 3, 4$ . By using Hausdorff distance(2.1), for  $j = 1, 2, 3$ , we obtain:

$$\begin{aligned} (s_0^-)^{(4)}(jh, r) &= f^-(jh, s_0(jh, r)), \\ (s_0^+)^{(4)}(jh, r) &= f^+(jh, s_0(jh, r)) \end{aligned} \quad (3.7)$$

by solving system(3.7), the value of  $\alpha_{i,0}$  for  $i = 5, 6, 7, 8$  are obtained and  $s_0(t)$  is constructed.

**Step Two:** The approximate solution  $s(t)$  in interval  $[kH, (k+1)H]$  for  $k = 1, \dots, n-1$  is obtained as follows:

$$s(t) = \sum_{i=0}^4 s_{4k}^{(i)}(t) \odot \frac{(t-4kh)^i}{i!} \oplus \sum_{i=5}^8 \alpha_{i,k} \odot \frac{(t-4kh)^i}{i!}, \quad (3.8)$$

where  $s_0(t)$  is obtained by step 1. The value of  $\alpha_{i,k}$  are to be determined so that  $s(t)$  satisfy the relations:

$$s^{(4)}(jh) = f(jh, s(jh)). \quad (3.9)$$

This means for  $j = 4k+1, 4k+2, 4k+3, 4k+4; k = 1, \dots, n-1$ ,



$$\begin{aligned}
(s^-)^{(4)}(jh, r) &= f^-(jh, s(jh, r)), \\
(s^+)^{(4)}(jh, r) &= f^+(jh, s(jh, r)),
\end{aligned} \tag{3.10}$$

by solving system (3.10), the values of  $\alpha_{i,k}$  are obtained.

Therefore the approximate solution is obtained as follows

$$s(t) = \begin{cases} s_{0,1}(t) & 0 \leq t \leq H, \\ \sum_{i=0}^4 s_{4k}^{(i)}(t) \odot \frac{(t-4kh)^i}{i!} \oplus \sum_{i=5}^8 \alpha_{i,k} \odot \frac{(t-4kh)^i}{i!}, & kH \leq t \leq (k+1)H \end{cases} \tag{3.11}$$

if  $y(t)$  is  $[(i) - gH]$ -differentiable and

$$s(t) = \begin{cases} s_{0,2}(t) & 0 \leq t \leq H, \\ \sum_{i=0}^4 s_{4k}^{(i)}(t) \odot \frac{(t-4kh)^i}{i!} \oplus \sum_{i=5}^8 \alpha_{i,k} \odot \frac{(t-4kh)^i}{i!}, & kH \leq t \leq (k+1)H \end{cases} \tag{3.12}$$

if  $y(t)$  is  $[(ii) - gH]$ -differentiable, and

$$s(t) = \begin{cases} s_{0,3}(t) & 0 \leq t \leq H, \\ \sum_{i=0}^4 s_{4k}^{(i)}(t) \odot \frac{(t-4kh)^i}{i!} \oplus \sum_{i=5}^8 \alpha_{i,k} \odot \frac{(t-4kh)^i}{i!}, & kH \leq t \leq (k+1)H \end{cases} \tag{3.13}$$

if  $y^{(i)}(t)$  are  $[(ii) - gH]$ -differentiable for  $i = 0, 2$ , and  $y^{(i)}(t)$  are  $[(i) - gH]$ -differentiable for  $i = 1, 3$ .

#### 4 Existence and uniqueness

In this section we prove that there exist a unique piecewise approximation solution  $s(t)$  where approximating the solution of fourth order fuzzy differential equation (3.1), provided that the size of the subinterval  $h$  satisfies some constraints.

**Theorem 4.1** *If  $h = \min\{h_1, h_2, h_3, h_4\}$ , where*

$$h_1 < \sqrt[4]{\frac{5}{L}}, \quad h_2 < \sqrt[4]{\frac{0.756}{L}}, \quad h_3 < \sqrt[4]{\frac{0.467}{L}}, \quad h_4 < \sqrt[4]{\frac{0.988}{L}} \quad (4.1)$$

then the approximate solution defined by (3.11) or (3.12), exists and unique.

**Proof:** Let  $t = jh$  and  $j = 3k + \eta$  for  $\eta = 1, 2, 3$ , therefore

$$s^{(4)}((4k + \eta)h) = s_{4k+\eta}^{(4)} = s_{4k}^{(4)} + \sum_{i=5}^8 \alpha_{i,k} \frac{(\eta h)^{i-4}}{(i-4)!} \quad (4.2)$$

where  $\eta = 1, \dots, 4$ . By solving system (4.2) we obtain:

$$\begin{aligned} \alpha_{5,k}^+ &= \frac{1}{12h} (48(s_{4k+1}^+)^{(4)} - 36(s_{4k+2}^+)^{(4)} \\ &\quad + 16(s_{4k+3}^+)^{(4)} - 3(s_{4k+4}^+)^{(4)} - 25(s_{4k}^+)^{(4)}), \end{aligned} \quad (4.3)$$

$$\begin{aligned} \alpha_{5,k}^- &= \frac{1}{12h} (48(s_{4k+1}^-)^{(4)} - 36(s_{4k+2}^-)^{(4)} \\ &\quad + 16(s_{4k+3}^-)^{(4)} - 3(s_{4k+4}^-)^{(4)} - 25(s_{4k}^-)^{(4)}), \end{aligned} \quad (4.4)$$

$$\begin{aligned} \alpha_{6,k}^+ &= \frac{-1}{12h^2} (104(s_{4k+1}^+)^{(4)} - 114(s_{4k+2}^+)^{(4)} \\ &\quad + 56(s_{4k+3}^+)^{(4)} - 11(s_{4k+4}^+)^{(4)} - 35(s_{4k}^+)^{(4)}), \end{aligned} \quad (4.5)$$

$$\begin{aligned} \alpha_{6,k}^- &= \frac{-1}{12h^2} (104(s_{4k+1}^-)^{(4)} - 114(s_{4k+2}^-)^{(4)} \\ &\quad + 56(s_{4k+3}^-)^{(4)} - 11(s_{4k+4}^-)^{(4)} - 35(s_{4k}^-)^{(4)}), \end{aligned} \quad (4.6)$$

$$\begin{aligned}\alpha_{7,k}^+ &= \frac{1}{2h^3}(18(s_{4k+1}^+)^{(4)} - 24(s_{4k+2}^+)^{(4)}) \\ &\quad + 14(s_{4k+3}^+)^{(4)} - 3(s_{4k+4}^+)^{(4)} - 5(s_{4k}^+)^{(4)},\end{aligned}\tag{4.7}$$

$$\begin{aligned}\alpha_{7,k}^- &= \frac{1}{2h^3}(18(s_{4k+1}^-)^{(4)} - 24(s_{4k+2}^-)^{(4)}) \\ &\quad + 14(s_{4k+3}^-)^{(4)} - 3(s_{4k+4}^-)^{(4)} - 5(s_{4k}^-)^{(4)},\end{aligned}\tag{4.8}$$

$$\begin{aligned}\alpha_{8,k}^+ &= \frac{-1}{h^4}(4(s_{4k+1}^+)^{(4)} - 6(s_{4k+2}^+)^{(4)}) \\ &\quad + 4(s_{4k+3}^+)^{(4)} - (s_{4k+4}^+)^{(4)} - (s_{4k}^+)^{(4)},\end{aligned}\tag{4.9}$$

$$\begin{aligned}\alpha_{8,k}^- &= \frac{-1}{h^4}(4(s_{4k+1}^-)^{(4)} - 6(s_{4k+2}^-)^{(4)}) \\ &\quad + 4(s_{4k+3}^-)^{(4)} - (s_{4k+4}^-)^{(4)} - (s_{4k}^-)^{(4)},\end{aligned}\tag{4.10}$$

To prove the existence and uniqueness of  $s(t)$ , let us define the operator  $G_k : \mathbb{R}_{\mathcal{F}} \rightarrow \mathbb{R}_{\mathcal{F}}$  by  $\alpha_{j,k} = G_v(\alpha_{j,k})$  for  $j = 5, 6, 7, 8$  and  $v = 1, 2, 3, 4$ . According to condition (3.2) and equations (4.3),(4.5),(4.7), (4.9) and (4.4),(4.6),(4.8), (4.10) we conclude that

$$\begin{aligned}D(G_1(\alpha_{5,k}) , G_1(\alpha_{5,k}^*)) &\tag{4.11} \\ &\leq L \frac{h^4}{12.5!} |48 - 36(2^5) + 16(3^5) - 3(4^5)| D(\alpha_{5,k}, \alpha_{5,k}^*),\end{aligned}$$

$$\begin{aligned}D(G_2(\alpha_{6,k}) , G_2(\alpha_{6,k}^*)) &\tag{4.12} \\ &\leq L \frac{h^4}{12.6!} |104 - 114(2^6) + 56(3^6) - 11(4^6)| D(\alpha_{6,k}, \alpha_{6,k}^*),\end{aligned}$$

$$\begin{aligned}
D(G_3(\alpha_{7,k}) , G_3(\alpha_{7,k}^*)) & \tag{4.13} \\
& \leq L \frac{h^4}{2.7!} |18 - 24(2^7) + 14(3^7) - 3(4^7)| D(\alpha_{7,k}, \alpha_{7,k}^*),
\end{aligned}$$

$$\begin{aligned}
D(G_4(\alpha_{8,k}) , G_4(\alpha_{8,k}^*)) & \tag{4.14} \\
& \leq L \frac{h^8}{8!} |4 - 6(2^8) + 4(3^8) - 4^8| D(\alpha_{8,k}, \alpha_{8,k}^*),
\end{aligned}$$

From Equations (4.11), (4.12), (4.13), (4.14), and

$$h_1 < \sqrt[4]{\frac{5}{L}}, \quad h_2 < \sqrt[4]{\frac{0.756}{L}}, \quad h_3 < \sqrt[4]{\frac{0.467}{L}}, \quad h_4 < \sqrt[4]{\frac{0.988}{L}}$$

it follows that  $G_v$ ,  $v = 1, 2, 3, 4$  are contraction operators. This implies the existence and uniqueness of approximate method under the stated conditions of theorem.

## 5 Numerical Example

**Example 5.1** Consider the fuzzy initial value problem

$$\begin{aligned}
y^{(4)}(t) &= y(t), \quad t \in [0, 1], \\
y(0) &= y'(0) = y'''(0) = y^{(3)}(0) = (r - 1, 1 - r),
\end{aligned}$$

$y(t)$  is  $[(i) - gH]$ -differentiable and the real solution is:

$$\begin{aligned}
y^-(t, r) &= (r - 1)e^t, \\
y^+(t, r) &= (1 - r)e^t.
\end{aligned}$$

We consider  $I_k = [kH, (k + 1)H]$ , for  $k = 0, 1$ ,  $H = 4h$  and  $h = 0.125$ .  $s_0(t)$ ,  $s_4(t)$  are obtained as follows:

$$\begin{aligned}
s_0^-(t) &= (r-1) + t(r-1) + \frac{t^2}{2}(r-1) + \frac{t^3}{3!}(r-1) + \frac{t^4}{4!}(r-1) \\
&+ \frac{t^5}{5!}(-.9999397099 + .9999397099r) + \frac{t^6}{6!}(-1.001990135 + 1.001990135r) \\
&+ \frac{t^7}{7!}(-.9672052893 + .9672052893r) + \frac{t^8}{8!}(-1.287374418 + 1.287374418r),
\end{aligned}$$

$$\begin{aligned}
s_0^+(t) &= (1-r) + t(1-r) + \frac{t^2}{2}(1-r) + \frac{t^3}{3!}(1-r) + \frac{t^4}{4!}(1-r) \\
&+ \frac{t^5}{5!}(.9999397099 - .9999397099r) + \frac{t^6}{6!}(1.001990135 - 1.001990135r) \\
&+ \frac{t^7}{7!}(.9672052893 - .9672052893r) + \frac{t^8}{8!}(1.287374418 - 1.287374418r),
\end{aligned}$$

$$\begin{aligned}
s_4^-(t) &= 1.648721270r - 1.648721270 \\
&+ (k-0.5)(1.648721270r - 1.648721270) \\
&+ \frac{(k-0.5)^2}{2}(1.648721270r - 1.648721270) \\
&+ \frac{(k-0.5)^3}{3!}(1.648721270r - 1.648721270) \\
&+ \frac{(k-0.5)^4}{4!}(1.648721270r - 1.648721270) \\
&+ \frac{(k-0.5)^5}{5!}(-1.648621871 + 1.648621871r) \\
&+ \frac{(k-0.5)^6}{6!}(-1.652002422 + 1.652002422r) \\
&+ \frac{(k-0.5)^7}{7!}(-1.594651978 + 1.594651978r) \\
&+ \frac{(k-0.5)^8}{8!}(-2.122522161 + 2.122522161r),
\end{aligned}$$

$$\begin{aligned}
s_4^+(t) &= 1.648721270 - 1.648721270r \\
&+ (k - 0.5)(1.648721270 - 1.648721270r) \\
&+ \frac{(k - 0.5)^2}{2}(1.648721270 - 1.648721270r) \\
&+ \frac{(k - 0.5)^3}{3!}(1.648721270 - 1.648721270r) \\
&+ \frac{(k - 0.5)^4}{4!}(1.648721270 - 1.648721270r) \\
&+ \frac{(k - 0.5)^5}{5!}(1.648621871 - 1.648621871r) \\
&+ \frac{(k - 0.5)^6}{6!}(1.652002422 - 1.652002422r) \\
&+ \frac{(k - 0.5)^7}{7!}(1.594651978 - 1.594651978r) \\
&+ \frac{(k - 0.5)^8}{8!}(2.122522161 - 2.122522161r),
\end{aligned}$$

Table 1  
Error of proposed method by Hausdorff distance in example 5.1

t	Error of Proposed method
0	0
0.1	0
0.2	0
0.3	0
0.4	$0.1 \times 10^{-8}$
0.5	$0.1 \times 10^{-8}$
0.6	$0.1 \times 10^{-8}$
0.7	$0.1 \times 10^{-8}$
0.8	$0.1 \times 10^{-8}$
0.9	$0.3 \times 10^{-8}$




Fig. 1. Approximate solution for example 5.1. Red points:  $s_0(t)$ ; green points:  $s_4(t)$ .

The approximated solution  $s(t)$ , for  $i = 0, 1$ , is plotted in Fig 1.

## 6 Conclusion

In this paper a new numerical method for solving forth order fuzzy differential equations under generalized differentiability was proposed. We used piecewise fuzzy polynomial of degree 8 based on the taylor expansion for approximating solutions of forth order fuzzy differential equations. Also, we can extend this method for  $N$ -th order fuzzy differential equations under generalized differentiability.

## References

- [1] G. Roussy, J. A. Percy, *Foundations and industrial applications of microwaves and radio frequency fields*, John Wiley, New York, 1995.
- [2] R. A. Van Gorder, K. Vajravelu, A variational formulation of the Nagumo reaction-diffusion equation and the Nagumo telegraph equation, *Nonlinear Analysis: Real World Applications* 4 (2010), 2957–2962.
- [3] L.A. Zadeh, The concept of a linguistic variable and its application to approximate reasoning, *Inform. Sci* 8 (1975), 199–249.
- [4] S. Abbasbandy, T. Allahviranloo, Numerical Solutions of Fuzzy Differential Equations By Taylor Method, *Journal of Computational Methods in Applied Mathematics* 2 (2002), 113–124.
- [5] T. Allahviranloo, N.Ahmady, E.Ahmady, Numerical solution of fuzzy differential equations by Predictor-Corrector method, *Information Sciences* 177/7 (2007), 1633–1647.
- [6] T. Allahviranloo, E.Ahmady, N.Ahmady, Nth-order fuzzy linear differential equations, *Information Sciences* 178 (2008), 1309–1324.
- [7] T. Allahviranloo, E.Ahmady, N.Ahmady, A method for solving nth order fuzzy linear differential equations, *International Journal of Computer Mathematics* (2009), 730 -742. DOI: 10.1080/00207160701704564
- [8] T. Allahviranloo, S. Abbasbandy, N. Ahmady and E. Ahmady, Improved predictor-corrector method for solving fuzzy initial value problems, *Information Sciences* 179 (2009), 945–955.
- [9] T. Allahviranloo, Z. Gouyandeh, A. Armand, A full fuzzy method for solving differential equation based on Taylor expansion, *Intelligent and Fuzzy Systems*, *Intelligent and Fuzzy Systems* 29 (2015), 1039-1055 DOI:10.3233/IFS-151713.
- [10] M.R. Balooch Shahryari, S. Salahshour, Improved predictor-corrector method for solving fuzzy differential equations under generalized differentiability, *Fuzzy Set Value Analysis 2012* (2012), 1–16.
- [11] B. Bede, I. J.Rudas, A. L.Bencsik, First order linear fuzzy differential equations under generalized differentiability, *Information Sciences* 177 (2007), 1648–1662.



- [12] B. Bede and S. G. Gal, Generalizations of the differentiability of fuzzy-number-valued functions with applications to fuzzy differential equations, *Fuzzy Set and Systems* 151 (2005), 581–599.
- [13] B. Bede and S. G. Gal, Remark on the new solutions of fuzzy differential equations *Chaos Solitons Fractals* (2006).
- [14] B. Bede and L. Stefanini, Solution of Fuzzy Differential Equations with generalized differentiability using LU-parametric representation *EUSFLAT1*(2011)785–790.
- [15] B. Bede and L. Stefanini, Generalized differentiability of fuzzy-valued functions *Fuzzy Sets and Systems* 230 (2013)119–141.
- [16] J. J. Buckley and T. Feuring, Fuzzy initial value problem for Nth-order linear differential equations *Fuzzy Sets and System* 121 (2001) 247-255.
- [17] S.L. Chang, L.A. Zadeh, On fuzzy mapping and control, *IEEE Trans, Systems Man Cybernet* 2 (1972) 30-34.
- [18] D. Dubois, H. Prade, Towards fuzzy differential calculus: Part 3, differentiation, *Fuzzy Sets and Systems* 8 (1982) 225–233.
- [19] D. Dubois, H. Prade, *Fundamentals of fuzzy sets*, Kluwer Academic Publishers, USA, (2000).
- [20] M. Friedman, M. Ma, A. Kandel, Numerical solutions of fuzzy differential and integral equations, *Fuzzy Sets and Systems* 106 (1999) 35-48.
- [21] D. N. Georgiou, J. J. Nieto, and R. Rodriguez-Lopez, Initial value problems for higher-order fuzzy differential equations, *Nonlinear Analysis: Theory, Methods and Applications* 63 (2005) 587–600.
- [22] O. Kaleva, Fuzzy differential equations. *Fuzzy Sets and Seystem* 24 (1987) 319-330.
- [23] New Results on Multiple solution for Nth-Order Fuzzy Differential Equations under Generalized Differentiability, *Boundary Value Problems* 2009 (2009) 1-13.
- [24] M. Ma, M. Friedman, A. Kandel, Numerical Solutions of fuzzy differential equations, *Fuzzy Sets and Systems* 105 (1999) 133-138.
- [25] S. Sallam, H.M. El-Hawary, A deficient spline function approximation to system of first order differential equations, *App. Math. Modelling* 7 (1983)380-382.

- [26] S. Seikkala, On the fuzzy initial value problem, *Fuzzy Sets and Systems* 24 (1987) 319-330.