



ω_0 –Nearest Points and ω_0 –Farthest Points in Normed Linear Spaces

H. Mazaheri ^{a,*}

^a*Faculty of Mathematics, Yazd University, Yazd, Iran*

Received 17 May 2015; accepted 5 January 2016

Abstract

In this paper we obtain a necessary and a sufficient condition for the set of ω_0 -nearest points (ω_0 -farthest points) to be non-empty or a singleton set in normed linear spaces. We shall find a necessary and a sufficient condition for a uniquely remotal set to be a singleton set.

Key words: Proximinal sets, Chebyshev sets, Farthest points, Uniquely remotal sets, Remotal sets, ω_0 -Nearest point, ω_0 -Farthest point.

2010 AMS Mathematics Subject Classification : 41A50, 41A52, 41A65.

* Corresponding author's E-mail:hmazaheri@yazd.ac.ir(H. Mazaheri)

1 Introduction

Franchetti and Singer [3] obtained some results on the characterization and existence of farthest points in normed linear spaces in terms of bounded linear functionals.

Let W be a non-empty subset of a normed linear space X . For any $x \in X$, the (possibly empty) set of best approximation x from W is defined by

$$P_W(x) = \{y \in W : \|x - y\| = d(x, W)\},$$

where $d(x, W) = \inf\{\|x - y\| : y \in W\}$.

For $\omega_0 \in W$, we have

$$P_W^{-1}(\omega_0) = \{x \in X : \omega_0 \in P_W(x)\} = \{x \in X : \|x - \omega_0\| = d(x, W)\},$$

which is called ω_0 -nearest points set. It is clear that, if $0, \omega_0 \in W$,

$$P_W^{-1}(\omega_0) = \omega_0 + P_W^{-1}(0).$$

Note that if $z \in P_W^{-1}(0)$, then $\alpha z \in P_{\alpha W}^{-1}(0)$ for every scalar α .

The subset W is said to be proximal if the set $P_W(x)$ is non-empty for every $x \in X$ and the set W is Chebyshev if $P_W(x)$ is a singleton set for every $x \in X$ (see [2-3, 9-10]).

Let W be a non-empty bounded subset of a real normed linear space X and $x \in X$. An element $g_0 \in W$ is called a farthest point to x in W if

$$\|g_0 - x\| = \rho(x, W) = \sup_{g \in W} \|g - x\|,$$

the (possibly empty) set of farthest points x from W is defined by

$$F_W(x) = \{y \in W : \|y - x\| = \rho(x, W)\}.$$

For $\omega_0 \in W$, we define

$$F_W^{-1}(\omega_0) = \{x \in X : \omega_0 \in F_W(x)\} = \{x \in X : \|x - \omega_0\| = \rho(x, W)\},$$

which is called ω_0 -farthest points set. It is clear that if $0, \omega_0] \in W$,

$$F_W^{-1}(\omega_0) = \omega_0 + F_W^{-1}(0).$$

Note that if $z \in F_W^{-1}(0)$, then $\alpha z \in F_{\alpha W}^{-1}(0)$ for every scalar α .

It is clear that for $x \in X$,

$$F_W(x) = (x - F_W^{-1}(0)) \cap W.$$

Let W be a bounded set in a normed linear space X . The set W is said to be remotal if the set $F_W(x)$ is non-empty for each $x \in X$, uniquely remotal if the set $F_W(x)$ consist of exactly one element for each $x \in X$,. (see [4, 6-8])

We will use the well-known fact about proximity.

Lemma 1 [9] *Let X be a normed linear space and W be a linear subspace of X . If W is proximal for $r > 0$, then there exists a $z \in P_W^{-1}(0)$.*

Lemma 2 [7] *Let W be a uniquely remotal subset of a normed space $(X, \|\cdot\|)$. Then a necessary and sufficient condition that W be singleton is that*

$$\|x - q_W(x)\| = \|y - q_W(x)\| \implies q_W(x) = q_W(y).$$

2 ω_0 -Nearest point sets and ω_0 -Farthest point sets

In this section, we will consider, ω_0 -nearest point sets and ω_0 -farthest point sets and uniquely remotal sets in normed linear spaces.

Theorem 3 *Let X be a normed linear space,*

(i) If W is a non-empty proximal subset of X and $P_W^{-1}(\omega_0)$ is singleton, then W is Chebyshev.

(ii) If W is a non-empty remotal bounded subset of X , $\omega_0 \in W$ and $F_W^{-1}(\omega_0)$ is singleton. Then W is uniquely remotal set.

Proof. (i) Suppose $x \in X$ and $g_1, g_2 \in P_W(x)$. Then $x - g_i + \omega_0 \in P_W^{-1}(\omega_0)$, for every $i = 1, 2$. Therefore $g_1 = g_2$.

(ii) The proof is similar to proof of (i)

Theorem 4 Suppose X is a strictly convex Banach space. If W is a non-empty bounded subset of X and $\omega_0 \in W$. If $F_W^{-1}(\omega_0) \neq \emptyset$ then $P_W^{-1}(\omega_0) \neq \emptyset$.

Proof. Form [5], if $z \in W$ is a farthest point from an $x \in X$, then z is also a nearest point in W . Now suppose $x \in F_W^{-1}(\omega_0)$, then $\omega_0 \in F_W(x)$, it follows that for some a $z \in X$, we have $\omega_0 \in P_W(z)$, therefore $z \in P_W^{-1}(\omega_0)$.

Example 2.1 Let X be a normed space and $W = \{x \in X : \|x\| \leq 1\}$. For $x \in X$, it is trivial for every $x \in X$, $d(x, W) = |1 - \|x\||$, $\delta(x, W) = |1 + \|x\||$ and $x \in P_W(\frac{x}{\|x\|})$ and $-x \in F_W(\frac{x}{\|x\|})$. Also it is clear that $P_W^{-1}(0) = F_W^{-1}(0) = W$

We know that in a normed linear space X , a vector x is said to be Birkhoff orthogonal to a vector y if the inequality $\|x\| \leq \|x + \alpha y\|$ holds for any real number α .

Theorem 5 Let $(H, \langle \cdot, \cdot \rangle)$ be an inner product space, W is a subspace of H . Then $P_W^{-1}(0) = W^\perp$ and for $\omega_0 \in W$ we have $P_W^{-1}(\omega_0) = \omega_0 + W^\perp$.

Proof. It is clear that if X is normed linear space, W is a subspace of X and $w \in W$. Then for any $x \in P_W^{-1}(0)$ and $x \perp w$. Therefore $x \in W^\perp$. If $x \in W^\perp$, then $x \perp w$ for every $w \in W$. Therefore $\|x\| \leq \|x + \alpha w\|$ for $w \in W$. Since W is a subspace, we have $\|x\| = d(x, W)$ and $x \in P^{-1}(0)$. Therefore $P_W^{-1}(0) = W^\perp$.

Example 2.2 (i) Let $X = \mathbb{R}^2$ with Euclidean norm, and $W = \{(x, x) : x \in \mathbb{R}\}$ be a subspace. From Theorem 2.2, $F_W^{-1}(0) = W^\perp = \langle (-1, 1) \rangle$.

(ii) Let $X = \mathbb{R}^2$ with Euclidean norm, and $W = \{(x, x) : 0 \leq x \leq 1\}$. Then $F_W^{-1}(0) = \{(x, y) : x \geq 1\}$.

Theorem 6 Suppose X is a normed linear space. (i) If W is a non-empty subset of X and $\omega_0 \in W$. Then W is proximal if and only if $X = W + P_W^{-1}(\omega_0)$. (ii) If W is a non-empty bounded subset of X and $\omega_0 \in W$. Then W is remotal if and only if $X = W + F_W^{-1}(\omega_0)$.

Proof. It is clear.

Theorem 7 Let X be a normed linear space.

(i) W a subspace of X with codimension one, and there exists a $z \in P_W^{-1}(0)$ and $X = W \oplus \langle z \rangle$, (where \oplus means that the sum decomposition of each element $x \in E$ is unique), then W is proximal.

(ii) W a proximal subspace of X and $P_W^{-1}(0) = \langle z \rangle$. Then W is Chebyshev.

(iii) W a non-empty bounded subset of X , $0 \in W$ and W is remotal, then there exists a $z \in F_W^{-1}(0)$.

(iv) W a non-empty bounded subset of X , $0 \in W$. If there exists a $z \in F_W^{-1}(0)$ and $X = W \oplus \langle z \rangle$, (where \oplus means that the sum decomposition of each element $x \in X$ is unique.) and $W = \beta W$ for every scalar β , then W is remotal.

(v) W a non-empty bounded subset of X , $0 \in W$ and there exists a unique $z \in X$ such that $F_W^{-1}(0) = \{z\}$. Then W is uniquely remotal

(vi) X a reflexive space and has the Kadec-Klee property. For every non-empty bounded subset of W of X and $0 \in W$, the set $F_W^{-1}(0)$

is compact.

Proof. (i) For arbitrary $x \in X \setminus W$, there exists a unique element $h \in W$ and the scalar α such that $x = h + \alpha z$. In this case $h \in P_W(x)$, and therefore W is proximal.

(ii) In this case, we show that $X = W \oplus P_W^{-1}(0)$. Since if $x \in X$, there exists a $g_0 \in P_W(x)$. Then $x = g_0 + (x - g_0)$ and $X = W + P_W^{-1}(0)$, also $W \cap P_W^{-1}(0) = \{0\}$. Now for $x \in X$, suppose $g_1, g_2 \in P_W(x)$. We have $x = g_1 + (x - g_1) = g_2 + (x - g_2)$ and the sum decomposition of each element $x \in X$ is unique. Therefore $g_1 = g_2$.

(iii) Suppose $x \in X \setminus W$, there exists a $g_0 \in F_W(X)$, then $z = x - g_0 \in F_W^{-1}(0)$.

(iv) For arbitrary $x \in X \setminus W$, there exists a unique element $h \in W$ and the scalar α such that $x = h + \alpha z$. In this case $h \in F_W(x)$, since $W = \alpha W$, therefore W is remotal.

(v) For $x \in X$, suppose $g_1, g_2 \in F_W(x)$, consider $z_i = x - g_i$. Therefore $z_i \in F_W^{-1}(0)$, for $i = 1, 2$, therefore $z_1 = z_2 = z$, and it follows that $g_1 = g_2$.

(vi) Since X is reflexive, the closed unit ball B_X is weakly compact. Consider the sequence $\{x_n\} \subseteq F_W^{-1}(0)$. We define $y_n = \frac{x_n}{\rho(x_n, W)}$. Therefore $y_n \in B_X$. Therefore there exists a subsequence $\{y_{n_k}\}$ and $y_0 \in B_X$ such that $y_{n_k} \rightharpoonup y_0$. Since X has Kadec-Klee property, $y_{n_k} \rightarrow y_0$. Also the sequence $\{\rho(x_n, W)\}$ is a bounded sequence and has a convergence subsequence $\{\rho(x_{n_l}, W)\}$ to k . Thus $x_{n_p} \rightarrow y_0 k$. Then the set $F_W^{-1}(0)$ is compact.

Theorem 8 Suppose X is a normed linear space.

(i) If W is a non-empty subspace of X and $\omega_0 \in W$. We have $P_W^{-1}(\omega_0) = X$ if and only if W is singleton and $W = \{\omega_0\}$.

(ii) If W is a non-empty bounded subset of X and $\omega_0 \in W$. We have $F_W^{-1}(\omega_0) = X$ if and only if W is singleton and $W = \{\omega_0\}$.

Proof. (i) If $P_W^{-1}(\omega_0) = X$, then for every $w \in W$, we have $w \in P_W^{-1}(\omega_0)$. Therefore $w = \omega_0$ and $W = \{\omega_0\}$. If $W = \{\omega_0\}$ and $x \in X$, then $d(x, W) = \|x - \omega_0\|$. Therefore $x \in P_W^{-1}(\omega_0)$.

(ii) Suppose $W = \{\omega_0\}$, if $x \in X$, then $q_x = \omega_0$. That is $\|x - \omega_0\| = \rho(x, W)$, therefore $x \in F_W^{-1}(\omega_0)$. It follows that $X = F_W^{-1}(\omega_0)$. If $X = F_W^{-1}(\omega_0)$, it is clear that W is remotal. Also, if for $x \in X$, there exist $w_1, w_2 \in F_W(x)$, then $w_1 = w_2 = \omega_0$. Therefore W is uniquely remotal. From Lemma 1.2, W is singleton. Therefore $W = \{\omega_0\}$.

Theorem 9 *Suppose X is a normed linear space. If W is a non-empty proximal subspace of X and $\omega_0 \in W$. If $P_W^{-1}(\omega_0)$ is convex. Then W is Chebyshev.*

Proof. Since $P_W^{-1}(\omega_0)$ is convex. The set $P_W^{-1}(0) = P_W^{-1}(\omega_0) - \omega_0$ is convex. From [5], W is Chebyshev.

Theorem 10 *Let $W \subseteq X$ be a proximal hyperplane and $\omega_0 \in W$. If $P_W(x)$ is compact for each $x \in X$. Then every sequence $\{x_n\}_{n \geq 1} \subseteq S_X$ with $x_n \in P_W^{-1}(\omega_0)$ for each n has a convergent subsequence.*

Proof. Suppose $P_W(x)$ is compact for each $x \in X$. If the sequence $\{x_n\}_{n \geq 1} \subseteq S_X$ with $x_n \in P_W^{-1}(\omega_0)$ for each n . Put $y_n = x_n - \omega_0 \in P_W^{-1}(0)$ and $\|y_n\| = d(x_n, W) = \|x_n\| = 1$. From [Theorem 2.1 ,6, 8], the sequences $\{y_n\}$ and $\{x_n\}$ has a convergent subsequence.

Theorem 11 *Let $W \subseteq X$ be a proximal hyperplane. If every sequence $\{x_n\}_{n \geq 1} \subseteq S_X$ with $x_n \in P_W^{-1}(0)$ for each n has a convergent subsequence. Then $P_W(x)$ is compact for each $x \in X$.*

Proof. If every sequence $\{x_n\}_{n \geq 1} \subseteq S_X$ with $x_n \in P_W^{-1}(0)$ for each n has a convergent subsequence. From [Theorem 2.1, 6, 8], W is quasi-Chebyshev subspace. It follows that $P_W(x)$ is compact for each $x \in X$.

Let W be a subspace of a normed space X . We define the quotient

space X/W to be the set of all sets $x + W$ of W together with the following operations:

$$(x + W) + (y + W) = (x + y) + W,$$

and

$$\lambda(x + W) = \lambda x + W,$$

for all $x, y \in X$ and arbitrary scalar λ . Then, the quotient space X/W is a normed space with the norm $\|x + W\| = \inf_{w \in W} \|x - w\|$.

Theorem 12 *Let M be a proximal subspace of a normed space X , W a proximal subspace of X containing M . If $P_W^{-1}(0) = \langle z \rangle$ for $z \in M$, then W/M is Chebyshev .*

Proof. From Lemma [2], W/M is proximal. Suppose there exist $z_1 + M, z_2 + M \in P_{W/M}(x + M)$. Since M is proximal there exist $m_1, m_2 \in M$ such that $d(z_1, M) = \|z_1 - m_1\|$ and $d(z_2, M) = \|z_2 - m_2\|$. Therefore $z_1 - m_1, z_2 - m_2 \in P_W^{-1}(0) = \langle z \rangle$, then $z_1 - m_1 = \alpha_1 z$ and $z_2 - m_2 = \alpha_2 z$. Therefore $z_1 + M = z_2 + M$.

For a Banach space X and closed subspace W of X , we denote its unit sphere by S_X . For $x \in X$ with $d(x, W) = 1$, let $Q_W(x) = x - P_W(x)$. It is easy to see that $Q_W(x) = \{z \in S_X : f(z) = f(x) \ \forall f \in W^\perp\}$.

For $f \in X^*$. we define the pre-duality map of X by

$$J_X(f) = \{z \in S_X : f(z) = \|f\|\}$$

.

Definition 2.1 [10] *Let X be a normed space, W be a subspace of X . Then W is a ω -Chebyshev subspace, if for every $x \in X$, $x + (P_W^{-1}(0) \cap S_X)$ is a nonempty and weakly compact set in X .*

Definition 2.2 [10] *A subspace W of a normed space X is called ω -boundedly compact if for every bounded sequence $\{y_n\}$ in W , there exists $x_0 \in W$ and a subsequence $\{y_{n_k}\}$ such that $y_{n_k} \rightharpoonup x_0$.*

In normed space X , suppose the unite sphere with center $\omega_0 \in X$ denoted by

$$S(\omega_0, 1) = \{x \in X : \|x - \omega_0\| = 1\}.$$

Theorem 13 *Let X be a normed space, W be a subspace of X and $\omega_0 \in X$. If $x \in X$ and the set $x + (P_W^{-1}(\omega_0) \cap S(\omega_0, 1))$ is nonempty and weakly compact, then W is ω -Chebyshev.*

Proof. It is clear, because

$$x + (P_W^{-1}(\omega_0) \cap S(\omega_0, 1)) = x + \omega_0 + (P_W^{-1}(0) \cap S_X).$$

Theorem 14 *Let X be a normed space, W be a subspace of X , $\omega_0 \in W$ and $\text{codim}W = 1$. Then the following statement are equivalent:*

- (i) $x + (P_W^{-1}(\omega_0) \cap S(\omega_0, 1))$ is nonempty and weakly compact.
- (ii) for every $f \in W^\perp$, $J_X(f)$ is weakly compact.
- (iii) for every $x \in X$, $P_W(x)$ is weakly compact.

Proof. (i) \Rightarrow (ii). Since W is ω -Chebyshev, from [Theorem 2.1, 10] for every $f \in W^\perp$, $J_X(f)$ is weakly compact.

(ii) \Rightarrow (iii). Theorem 2.1 of [10].

(iii) \Rightarrow (i). Suppose $x \in X$, and $\{z_n\} \subseteq x + (P_W^{-1}(\omega_0) \cap S(\omega_0, 1))$, then $\{z_n - \omega_0 \in x + (P_W^{-1}(0) \cap S_X)\}$. Therefore the sequence $\{z_n - \omega_0\}$ has a weakly convergent subsequence, it follows that $\{z_n\}$ has a weakly convergent subsequence. Also $x + \omega_0 \in X$ and W is ω -Chebyshev, therefore $x + (P_W^{-1}(\omega_0) \cap S(\omega_0, 1)) \neq \emptyset$.

Theorem 15 *Let X be a normed space, W be a subspace of X , $\omega_0 \in W$. If $P_W^{-1}(\omega_0)$ is ω -boundedly compact. Then $P_W(x)$ is weakly compact for every $x \in X$.*

Proof. Because $P_W^{-1}(0) = P_W^{-1}(\omega_0) - \omega_0$ is ω -boundedly compact, it

follows that from [Theorem 2.3, 10] the set $P_W(x)$ is weakly compact for every $x \in X$.

Corollary 16 *Let X be a normed space, W be a subspace of X , $\omega_0 \in W$. Then $P_W^{-1}(\omega_0)$ is ω -boundedly compact if and only if W is ω -Chebyshev.*

Acknowledgements

This research project is supported by intelligent robust research center of yazd university. The authors are highly grateful to the referees for their valuable comments and suggestions for improving the paper.

References

- [1] G. C. Ahuja, T. D. Narang and Swaran Trehan, On farthest points, J.Indian Math. Soc. **39**(1975), 293-297.
- [2] E. W., Cheney, D. E., Wulbert, The existence and unicity of best approximations. Math. Scand. **24** (1969) 113-140.
- [3] J. Fletcher, W. B. Moors, Chebyshev sets. J. Aust. Math. Soc. **98** (2015), 161-231.
- [4] C. Franchetti and I. Singer, Deviation and farthest points in normed spaces, Rev. Roum Math. Pures et Appl, **24**(1979), 373-381.
- [5] H. Mazaheri and F. M., Maalek Ghaini, Quasi-orthogonality of the best approximant sets. Nonlinear Anal. **65** (2006), 534537.
- [6] H. Mohebi, On quasi-Chebyshev subspaces of Banach spaces. J. Approx. Theory 107 (2000), 87-95.
- [7] T. D. Narang, On farthest points in metric spaces. J. Korea Soc. Math. Educ. Ser. B Pure Appl. Math. **9** (2002), 1-7.
- [8] D. Narayana; T. S. S. R. Rao, Some remarks on quasi-Chebyshev subspaces. J. Math. Anal. Appl. 321 (2006), 193-197.

- [9] A. Niknam, On uniquely remotal sets. *Indian J. Pure Appl. Math.* 15 (1984), 1079-1083.
- [10] M. Shams; H. Mazaheri; S. M. Vaezpour, On w -Chebyshev subspaces in Banach spaces. *Bull. Korean Math. Soc.* 45 (2008), 601-606.
- [11] Sangeeta, T. D. Narang, On the farthest points in convex metric spaces and linear metric spaces. *Publ. Inst. Math. (Beograd) (N.S.)* **95** (109) (2014), 229-238.
- [12] I. Singer, *Best Approximation in Normed Linear Spaces by Elements of Linear Subspaces*, Springer-Verlag, New York, Heidelberg, Berlin, 1970.
- [13] I. Singer, *The Theory of Best Approximation and Functional Analysis*, CBMS Reg. Confer. Series in Applied Math. No.13, SIAM, Philadelphia, 1974.